

DETECTING THE ORIENTATION OF LONG LINKS BY FINITE TYPE INVARIANTS

S.V.DUZHIN, M.V.KAREV

ABSTRACT. We prove the existence of a degree 7 Vassiliev invariant of long (or string) two-component links which is not preserved under the simultaneous change of orientation of both components. The non-invertibility of this invariant can be detected by the standard weight system with values in the tensor square of the universal enveloping algebra for $\mathfrak{gl}(n)$.

1. INTRODUCTION


It is well known that classical knot polynomials and quantum invariants in general take equal values on a knot and its inverse. The class of Vassiliev invariants is strictly wider [Vog], and the problem whether they can be used to tell a knot from its inverse is open and seemingly difficult. In this paper, we discuss the corresponding problem for links with more than 1 component.

Let S_p^1 be the disjoint union of p numbered copies of an oriented circle, $\mathbb{R}_p^1$ the disjoint union of p numbered copies of an oriented real line, and I_p^1 the disjoint union of p numbered copies of an oriented interval $[0, 1]$.

Definition. *An p -component closed link is a smooth embedding of S_p^1 into oriented 3-space $\mathbb{R}^3$, considered up to a component-preserving isotopy.*

An p -component long link is a smooth embedding of $\mathbb{R}_p^1$ into oriented 3-space $\mathbb{R}^3$ with fixed behavior at infinity: $x_i(t) = [i, 0, t]$ for $|t| > C$, considered up to an isotopy, fixed at infinity, and a Euclidean movement of space.

An p -component string link is a smooth embedding of I_p^1 into the slice $\mathbb{R}^2 \times [0, 1] \subset \mathbb{R}^3$ with fixed endpoints: $x_i(t) = [i, 0, t]$ for $t = 0, 1$, considered up to an isotopy, fixed at the boundary, and a Euclidean movement of the slice.



It is immediately clear that the theories of long and string links are equivalent. In the case of knots ($p = 1$) they are also equivalent to the theory of closed links; this is not true when $p > 1$.

The *problem of invertibility* is stated as follows. Given a link L , let L' be the same link with the orientation of all components reversed. May it happen that L' is not equivalent to L ? If yes, then what invariants can be used to detect that difference?

For knots, the problem was standing for a long time, until Trotter [Tro] proved in 1964 that some knots (e.g. pretzel knot with parameters (3, 5, 7)) are non-invertible. The simplest non-invertible knot is 8_{17} ([Ka]). However, it is not known if the two knots 8_{17} and $8'_{17}$, or any other pair of mutually inverse knots, can be distinguished by finite type invariants.

In the case of links, the only published result is a theorem of X.-S. Lin [L2] that asserts the non-invertibility of Vassiliev invariants for closed links with at least 6 components. Other papers containing the words ‘links’, ‘Vassiliev invariants’, ‘invertibility’ (‘separation’, ‘detection’ etc) in their titles/abstracts have smaller relevance to the plain problem we are speaking about, e.g. D.Bar-Natan [BN2] studies *homotopy* string links invariants, X.-S. Lin [L1] uses a modified definition of finite type invariants, etc.

The present text deals with the problem of detecting the orientation by finite type invariants for *long links*. The reformulation of the problem in terms of chord diagrams shows immediately that the answer is positive for $p > 2$. In the case of 2-component links ($p = 2$) the problem is nontrivial; below, we give a proof of the following

Theorem. *There exists a Vassiliev invariant f of order no greater than 7 and a 2-component long link L such that $f(L') \neq f(L)$.*

We will give two proofs of this theorem, both based on direct calculations. The first proof (Proposition 1, page 5) uses chord diagrams and requires computer calculations. The second one (Proposition 2, page 8) uses Jacobi diagrams and can be done by hand. Frankly speaking, both of these proofs refer to the invariants of *framed* links. In Section 6 we explain why they also imply the same result in the unframed case.

2. REDUCTION TO CHORD DIAGRAMS

Finite type invariants for various types of links are defined in the same way as they are in the classical case of closed knots, see [BN1, Sta].

Let $\mathbb{F}$ be a field of characteristic 0, e.g. $\mathbb{Q}$ or $\mathbb{C}$. Denote by $V_n(p)$ the vector space of $\mathbb{F}$ -valued Vassiliev invariants for p -component long links of degree no greater than n , by $\mathcal{A}_n(p)$ the space spanned by chord diagrams of degree n on p strings modulo 4-term relations, and by $W_n(p) = \mathcal{A}_n^*(p) = \text{Hom}_{\mathbb{Q}}(\mathcal{A}_n(p), \mathbb{F})$ the corresponding space of *weight systems*. Then there is a linear map $\sigma_n^p : V_n(p) \rightarrow W_n(p)$ (taking the *symbol* of a Vassiliev invariant) whose kernel coincides with $V_{n-1}(p)$ and whose image consists of all weight systems that vanish on chord diagrams with isolated chords. For *framed* links, the image equals the entire $W_n(p)$, and, until Section 6, we will actually be speaking about invariants of framed links.


As before, we denote by L' the inverse of a link L . Setting $\tau_V(f)(L) = f(L')$ for $f \in V$, we obtain an involution τ_V in the space of Vassiliev invariants. On the level of chord diagrams, the corresponding operation τ_A acts as follows:

$$\tau_A : \begin{array}{c} \uparrow \quad \uparrow \\ | \quad | \\ \diagdown \quad \diagup \\ | \quad | \\ \diagup \quad \diagdown \\ | \quad | \\ \downarrow \quad \downarrow \end{array} \mapsto \begin{array}{c} | \quad | \\ \diagdown \quad \diagup \\ | \quad | \\ \diagup \quad \diagdown \\ | \quad | \\ \downarrow \quad \downarrow \end{array} = \begin{array}{c} \uparrow \quad \uparrow \\ | \quad | \\ \diagdown \quad \diagup \\ | \quad | \\ \diagup \quad \diagdown \\ | \quad | \\ \downarrow \quad \downarrow \end{array},$$

i.e. it changes the orientation of all the strings of a chord diagram, or, which is the same thing, reflects the plane picture of the diagram in a horizontal line, assuming that the support is drawn vertically (and preserves its orientation).


Instead of usual chord diagrams only, the same space $\mathcal{A}(p)$ may be spanned by all generalized chord diagrams (*Chinese character diagrams* in the terminology of [BN1]), i.e. 1-3-valent graphs, attached to the same support $(\mathbb{R}_p^1)$ with a cyclic order of edges specified in every 3-valent vertex.

Generalized chord diagrams can be understood as linear combinations of ordinary chord diagrams by iterative applications of the STU rule, e.g.



(here and below, we use the *blackboard convention*: edges around trivalent vertices for every diagram drawn on paper are always ordered counterclockwise).

For generalized diagrams, one should be more cautious with the definition of the involution τ_A . In fact, extending the previously defined map τ_A from chord diagrams to generalized diagrams by STU relations, one arrives at the following rule:




i.e. one must either change the orientation of all strings *and multiply the result by ± 1* depending on the parity of the number of trivalent vertices, or, again, simply reflect the plane picture of the diagram in a horizontal line.

Lemma 1. *The involution τ_A is the graded part of the involution τ_V , i.e. the following square commutes:*

$$\begin{array}{ccc} V_n(p) & \xrightarrow{\sigma_n^p} & W_n(p) \\ \downarrow \tau_V & & \downarrow \tau_A^* \\ V_n(p) & \xrightarrow{\sigma_n^p} & W_n(p) \end{array}$$

Proof. This follows immediately from the definition of σ ([BN1, CD]). $\square$

The problem of detecting the inversion of orientation on long links by finite type invariants is thus equivalent to the problem: *find a chord diagram on p lines which is different from its inverse modulo $4T$ relations.* If $p \geq 3$, then there is a straightforward example:




For $n = 2$ the problem is non-trivial, because, as one can check, chord diagrams of small degrees (like the one depicted on page 2) are all τ_A -invariant. In the next section, we indicate a diagram which is not.

3. FIRST PROOF OF THE THEOREM

To prove that a certain element of the space $\mathcal{A}(p)$ is non-zero, one can use *weight systems*, i.e. linear functionals on these spaces. A powerful weight system is provided by Kontsevich's homomorphism $\varphi = \varphi_{\mathfrak{g}} : \mathcal{A}(p) \rightarrow U(\mathfrak{g})^{\otimes p}$ for a metrized Lie algebra $\mathfrak{g}$ (see [Kon, CD]). In fact, φ takes values in the $\mathfrak{g}$ -invariant subalgebra $U(p) = [U(\mathfrak{g})^{\otimes p}]^{\mathfrak{g}}$. We give an explicit description for the case $\mathfrak{g} = \mathfrak{gl}_N$, using the basis e_{ij} (matrix with one 1 and many 0's) and the metric defined by the conjugacy rule $e_{ij}^* = e_{ji}$.

Lemma 2. *Let $D \in \mathcal{A}(p)$ be a (generalized) chord diagram on p strings. The element $\varphi(D) \in U(\mathfrak{gl}_N)^{\otimes p}$ can be obtained as follows. Consider the alternating sum of all resolutions [BN1] of the inner triple points of the diagram D . For each resolution label the connected components of the obtained diagram by different independent indices; then replace each pair of adjacent indices by e_{ij} and take the sum over all indices from 1 to N . (Closed components, if any, turn into factors N .)*

The proof repeats the proof for the particular case $p = 1$ that can be found in [BN1, CD]. We give an example instead:



Therefore, the image of this diagram under φ is

$$\sum_{i,j,k,l=1}^N (e_{ij}e_{jk} \otimes e_{li}e_{kl} - e_{ij}e_{kl} \otimes e_{li}e_{jk} - e_{ij}e_{kl} \otimes e_{jk}e_{li} + e_{ij}e_{ki} \otimes e_{jl}e_{lk}).$$

Note that the order of the factors e_{ij} agrees with the orientation of each component of the support, but the order of the two subscripts (i, j) in each factor follows a go-round pattern, in our case bottom-to-top for the left line and top-to-bottom for the right line.

Denote by τ_U the operator in $U(\mathfrak{gl}_N)^{\otimes p}$ that rewrites each monomial backwards and changes the order of the two subscripts in each generator e_{ij} , e.g. $\tau_U(e_{12}e_{23} \otimes e_{13}e_{24}) = e_{32}e_{21} \otimes e_{42}e_{31}$. This is an involution that preserves the ad-invariant subspace $U(p) = [U(\mathfrak{gl}_N)^{\otimes p}]^{\mathfrak{g}}$.

The construction of φ shows that the following assertion holds.


Lemma 3. *The square*

$$\begin{array}{ccc} \mathcal{A}(p) & \xrightarrow{\varphi} & U(p) \\ \downarrow \tau_A & & \downarrow \tau_U \\ \mathcal{A}(p) & \xrightarrow{\varphi} & U(p) \end{array}$$

is commutative.

It follows that the non-invertibility of a chord diagram can be checked on the level of the universal enveloping algebra: if the φ -image of a diagram is not τ_U -invariant, then the diagram itself is not invariant under τ_A . Therefore, the next proposition gives the first proof of our Theorem.

Proposition 1. *Each of the following diagrams is different from its image under the involution τ_A :*



Proof. A direct computer calculation of the φ -images of each diagram and its inverse shows that they are different for the Lie algebra $\mathfrak{gl}_4$ (and hence for any $\mathfrak{gl}_N$ with $n \geq 4$). The calculations are organized as follows: we fix a lexicographical ordering of the basic elements e_{ij} and transform the expression for $\varphi(D)$ obtained by the above algorithm using the commutator relations between the generators in order to rewrite each monomial lexicographically. The programs, input and output files are available at [Calc]. The result for the left-hand side diagram, for example, is a polynomial consisting of 58378 terms; its calculation requires several hours on a reasonably fast modern PC. $\square$

4. REDUCTION TO JACOBI DIAGRAMS

Another, yet simpler, reformulation of the invertibility problem, can be given in terms of colored *Jacobi* diagrams. A colored Jacobi diagram is the same thing as a *Chinese character* as defined in [BN1], only its univalent vertices are labeled by p colors. The *space* of colored Jacobi diagrams $\mathcal{B}(p)$ is defined as a vector space formally generated by all p -colored Jacobi diagrams modulo two kinds of relations: antisymmetry and IHX.

Similarly to Theorem 8 of [BN1], one can prove that there is a symmetrization map $\chi : \mathcal{B}(p) \rightarrow \mathcal{A}(p)$ which provides a linear isomorphism between the two spaces (see also [LM]).

The map χ is defined as follows — we explain that in the case $p = 2$. Let D be a Jacobi diagram with k ‘legs’ of color 1 and l ‘legs’ of color 2, then $\chi(D)$ is the average of all the $k!l!$ ways to attach 1-colored legs to the first line of the support and 2-colored legs to the second line of the support, e.g.

$$\begin{array}{c} \bullet^2 \\ | \\ \bullet^1 \end{array} \begin{array}{c} \bullet^2 \\ | \\ \bullet^1 \end{array} \mapsto \frac{1}{2} \left(\begin{array}{c} \uparrow \uparrow \\ \bullet \bullet \\ | | \\ \bullet \bullet \\ \downarrow \downarrow \\ 1 \quad 2 \end{array} + \begin{array}{c} \uparrow \uparrow \\ \bullet \bullet \\ | | \\ \bullet \bullet \\ \downarrow \downarrow \\ 1 \quad 2 \end{array} \right)$$

Colored Jacobi diagrams can thus be viewed as symmetric elements of the space $\mathcal{A}(p)$, much in the same way as commutative polynomials over a Lie algebra $\mathfrak{g}$ can be viewed as symmetric elements of the universal enveloping algebra $U(\mathfrak{g})$, by virtue of the PBW theorem.

The isomorphism χ is very important for our needs because the involution τ_A takes an especially simple form when transferred to $\mathcal{B}(p)$ via this isomorphism. In fact, the following assertion holds:

Lemma 4. *Let $\tau_B : \mathcal{B}(p) \rightarrow \mathcal{B}(p)$ be the linear operator defined as identity on each Jacobi diagram with an even number of legs and as multiplication by -1 on each*

Jacobi diagram with an odd number of legs. Then the square

$$\begin{array}{ccc} \mathcal{B}(p) & \xrightarrow{\chi} & \mathcal{A}(p) \\ \downarrow \tau_B & & \downarrow \tau_A \\ \mathcal{B}(p) & \xrightarrow{\chi} & \mathcal{A}(p) \end{array}$$

is commutative.

Proof. The proof smoothly follows from the definitions of both τ_A and τ_B given above. A simple illustration might be helpful to understand how it goes. Let

$$D = \text{diagram of a circle with two internal vertical lines and two external legs labeled 1 and 2.}$$

Then by definition

$$\tau_B(D) = - \text{diagram of a circle with two internal vertical lines and two external legs labeled 1 and 2, with a sign change.}$$

Now

$$\chi(D) = \frac{1}{2} \text{diagram 1} + \frac{1}{2} \text{diagram 2}$$

and hence

$$\tau_A(\chi(D)) = -\frac{1}{2} \text{diagram 3} - \frac{1}{2} \text{diagram 4} = -\frac{1}{2} \text{diagram 5} - \frac{1}{2} \text{diagram 6} = \chi(\tau_B(D)).$$

□

The problem of finite type invertibility for 2-component long links can now be restated as follows: *is there a non-zero 2-colored Jacobi diagram with an odd number of legs?* In the next section, we give an example of such a diagram.

5. SECOND PROOF OF THE THEOREM

Take a metrized Lie algebra $\mathfrak{g}$ and denote by $S(\mathfrak{g})$ the symmetric algebra of the vector space $\mathfrak{g}$. The weight system $\mathcal{B}(1) \rightarrow S(\mathfrak{g})$ described in [BN1, CD] has an immediate generalization to an arbitrary value of p giving a homomorphism $\psi : \mathcal{B}(p) \rightarrow S(\mathfrak{g})^{\otimes p}$ whose image lies in the $\mathfrak{g}$ -invariant subalgebra $S(p) = [S(\mathfrak{g})^{\otimes p}]^{\mathfrak{g}}$.

Lemma 5. *The mapping ψ fits into a commutative diagram*

$$\begin{array}{ccc} \mathcal{B}(p) & \xrightarrow{\psi} & S(p) \\ \downarrow \chi & & \downarrow \pi \\ \mathcal{A}(p) & \xrightarrow{\varphi} & U(p) \end{array}$$

where χ is the isomorphism defined in the previous section, φ is the Kontsevich weight system for the algebra $\mathcal{A}(p)$, and π is the Poincaré-Birkhoff-Witt isomorphism raised to the n -th tensor power.

Proof. The proof is straightforward, just like in the conventional case $p = 1$. $\square$

For the Lie algebra $\mathfrak{g} = \mathfrak{gl}_N$ there is a pictorial algorithm for the calculation of ψ , similar to the procedure for φ described in section 3. The only difference is that now the basic elements e_{ij} are commutative, so we do not have to bother about the order of factors in the monomials. Here is a simple example:

$$\begin{aligned} \psi : \quad & \begin{array}{c} 2 \\ | \\ 1 \quad \bullet \quad 2 \end{array} \mapsto \begin{array}{c} j \quad i \\ | \quad | \\ i \quad 1 \quad 2 \quad i \\ \diagdown \quad \diagup \\ k \quad k \end{array} - \begin{array}{c} j \quad i \\ | \quad | \\ k \quad 1 \quad 2 \quad k \\ \diagdown \quad \diagup \\ i \quad j \end{array} \\ & \mapsto \sum_{i,j,k=1}^N (e_{jk} \otimes e_{ij}e_{ki} - e_{ki} \otimes e_{ij}e_{jk}) = 0. \end{aligned}$$

A big advantage that the mapping ψ has over φ is that the $\mathfrak{g}$ -invariant part of $S(\mathfrak{gl}_N)^{\otimes p}$ has a rather transparent structure. A p -colored *necklace* of order n is a combinatorial object defined as a sequence of n numbers between 1 and p , considered up to cyclic shifts; it can be best viewed as a circular arrangement of n p -colored beads. To every necklace one can assign an element of $S(\mathfrak{gl}_N)^{\otimes p}$, which turns out to be ad-invariant, as follows. Sprinkle different indices over the arcs of the necklace; every bead transforms into e_{ij} where i is the index written on the incoming arc and j is the index of the outgoing arc; put this element e_{ij} into the tensor factor whose number is the color of the bead and take the sum over all indices from 1 to N . For example:

$$[1212] = \begin{array}{c} \bullet \quad \bullet \\ \curvearrowright \end{array} \mapsto \begin{array}{c} j \quad i \\ \curvearrowright \\ k \quad l \end{array} \mapsto \sum_{i,j,k,l=1}^N (e_{ij}e_{kl} \otimes e_{jk}e_{li}) =: x_{1212}.$$

We denote such *necklace elements* by x_μ where μ is the lexicographically smallest sequence corresponding to the given necklace.

Lemma 6. *For the Lie algebra $\mathfrak{g} = \mathfrak{gl}_N$ the $\mathfrak{g}$ -invariant part of $S(\mathfrak{g})^{\otimes p}$, as an algebra, is generated by the necklace elements. Algebraic relations between necklace elements of a given degree may exist for small values of N , but they disappear as $N \rightarrow \infty$.*

Proof. Statements close to this lemma are proved in [W] and [VP]. We are sure that the fact itself must be written up somewhere, and we would be grateful to anybody who can give us an exact reference. $\square$

In the algebra generated by necklaces there is an involution τ_S that inverts the orientation of each necklace. For $p = 2$, in small degrees, up to 5, it is identical; the minimal necklace which is not fixed by τ_S , is x_{112122} . It turns out that the operation τ_S agrees with all other inversions denoted by τ with various subscripts in this paper.

Lemma 7. *The involutions of the change of orientation in the spaces $\mathcal{A}(p)$, $\mathcal{B}(p)$, $S(p)$, $U(p)$ commute with the four arrows of the diagram in Lemma (5). More precisely, we have a commutative cube*


$$\begin{array}{ccccc}
 & \mathcal{B}(p) & \xrightarrow{\psi} & S(p) & \\
 \tau_B \swarrow & \downarrow & & \searrow \tau_S & \\
 \mathcal{B}(p) & \xrightarrow{\psi} & S(p) & & \\
 \downarrow \chi & \downarrow \chi & \downarrow \pi & \downarrow \pi & \\
 & \mathcal{A}(p) & \xrightarrow{\varphi} & U(p) & \\
 \tau_A \swarrow & \downarrow & & \searrow \tau_U & \\
 \mathcal{A}(p) & \xrightarrow{\varphi} & U(p) & &
 \end{array}$$

(recall that $\mathfrak{g} = \mathfrak{gl}_N$, $S(p) = [S(\mathfrak{g})^{\otimes p}]^{\mathfrak{g}}$, $U(p) = [U(\mathfrak{g})^{\otimes p}]^{\mathfrak{g}}$; the maps were all defined in the text above, in particular, π is the Poincaré-Birkhoff-Witt isomorphism).

Proof. What we are really interested in is the commutativity of the top face of the cube. Since all χ 's, π 's and τ 's are linear isomorphisms, this follows from the commutativity of the remaining 5 faces which was proved in various places of this paper above. $\square$

The commutativity of the upper face of the cube implies that, to prove the non- τ_B -invariance of a Jacobi diagram (an element of $\mathcal{B}(p)$) it is enough to prove the non- τ_S -invariance of its image under ψ in the necklace algebra. Since the minimal degree of a non-invertible necklace is 6 and we need to find a non-invertible diagram of an odd degree, the smallest example should be searched in degree 7. And, indeed, it exists — thus providing a second proof of the Theorem.

Proposition 2. *The following diagram is non-zero as an element of the space $B(2)$:*



Proof. A direct computation shows that the image of this diagram in the algebra $S(2)$ (for N big enough), when expressed through necklaces, equals

$$N(x_{1121222} - x_{1122212}) + 3x_2(x_{112212} - x_{112122}),$$

which is different from zero. The entire expression for $\psi(H)$ involves 128 terms, out of which only 8 contain nonsymmetric necklaces, e.g.

$$\mapsto -x_2 x_{112122}.$$

The same result can also be obtained by a computer program available at [Calc]. $\square$

6. DEFRAMING

As we mentioned before, the proofs given in Sections 3 and 5, actually refer to the case of *framed* links, because the 1-term relations were not taken into consideration. In principle, detecting the orientation of a framed link is easier, because it contains an additional structure which may not be preserved by the inversion. Note that an unframed link can be considered as a framed one, so the Theorem in the unframed case is in any case stronger. Here we explain why does it hold.

Indeed, let $\mathcal{A}'(2)$ be the space $\mathcal{A}(2)$ modded out by 1-term relations, i.e. by the ideal generated by the diagrams a_1 (with 1 chord on the first component of the support) and a_2 (with 1 chord on the second component of the support): $\mathcal{A}'(2) = \mathcal{A}(2)/\langle a_1, a_2 \rangle$, where angular brackets denote the 2-sided ideal with given generators. The quotient algebra $\mathcal{A}'(2)$ can also be considered as a subalgebra of $\mathcal{A}(2)$: by the structure theorem for cocommutative Hopf algebras, $\mathcal{A}(2)$ is the universal enveloping algebra over its primitive subspace P , so if we take the subspace of P spanned by all connected diagrams except for a_1 and a_2 , we will obtain an inclusion $\mathcal{A}'(2) \subset \mathcal{A}(2)$. Since the diagrams displayed in Proposition 1 belong to $\mathcal{A}'(2)$, this means that non-invertibility holds for unframed long links, too.

Now let $\chi_2 : \mathcal{A}(2) \rightarrow \mathcal{B}(2)$ be the isomorphism discussed above.

Lemma 8. *The subspace $\chi^{-1}(\mathcal{A}'(2)) = \mathcal{B}'(2)$ coincides with the subalgebra of $\mathcal{B}(2)$ generated by all connected Jacobi diagrams except $b_1 = \text{---}\overset{1}{\bullet}\text{---}\overset{1}{\bullet}$ and $b_2 = \text{---}\overset{2}{\bullet}\text{---}\overset{2}{\bullet}$.*

Proof. The assertion is non-trivial, because χ does not preserve the multiplication, and the subalgebras $\mathcal{A}'(2)$ and $\mathcal{B}'(2)$ are described by their generators in the sense of different multiplications. However, a little thinking about the action of the operator χ shows that its value on a product of Jacobi diagrams different from b_1 and b_2 is equal to a linear combination of products of connected chord diagrams different from a_1 and a_2 . $\square$

The lemma shows that the ‘heptapus’ element displayed in Proposition 2 belongs indeed to the subalgebra $\mathcal{B}'(2)$ responsible for the Vassiliev invariants of unframed 2-component long links. Since it is non-zero, the existence of a 7 order invariant that can distinguish the orientation follows.

7. RELATED OPEN PROBLEMS

Problem 1. Is there a τ -equivariant linear mapping $\mathcal{B}(1) \rightarrow \mathcal{B}(2)$ whose image is not a priori contained in the τ -invariant subspace of $\mathcal{B}(2)$?


Remark. If such a mapping existed, then one could try to apply the techniques of this paper to the invertibility problem for *knots*. The standard doubling operator $\Delta : \mathcal{B}(1) \rightarrow \mathcal{B}(2)$ (as in [BN1, LM]) is no good, as one can easily see.

Problem 2. Do Vassiliev invariants detect invertibility of closed 2-component links?

Remark. The combinatorial object responsible for finite type invariants of closed links is the space of diagrams on 2 circles, call it $\mathcal{A}(S_2^1)$. There is an evident epimorphism $\mathcal{A}(\mathbb{R}_2^1) \rightarrow \mathcal{A}(S_2^1)$, but its kernel is non-trivial, so it cannot be used to define a Kontsevich type weight system with values in $U^{\otimes 2}$.

Problem 3. Is 7 the minimal degree of Vassiliev invariants that can be used to distinguish the orientation of long links with 2 components?

Remark. The question is reduced to the existence of non-invertible diagrams of degree smaller than 7. For example, we do not know if the following diagram



is zero or not as an element of $\mathcal{A}_5(2)$.

ACKNOWLEDGEMENTS

We thank A. Khoroshkin, S. Loktev and E. B. Vinberg for useful comments on invariant theory, and S. Lando who read the text and made some useful remarks. The first author was supported by RFBR grant 05-01-01012a.

REFERENCES

- [BN1] D. Bar-Natan, *On the Vassiliev knot invariants*, Topology, **34** (1995) 423–472.
- [BN2] D. Bar-Natan, *Vassiliev Homotopy String Link Invariants*, Journal of Knot Theory and its Ramifications 4-1 (1995) 13–32.
- [Calc] S. Duzhin. *Computer programs and data files for the calculation of the weight systems φ and ψ* . Online at <http://www.pdmi.ras.ru/~arnsem/dataprogram/OrLinks/>.
- [CD] S. V. Chmutov and S. V. Duzhin, *CDBook. Introduction to Vassiliev Knot invariants*, preliminary draft version of a book, Online at <http://www.pdmi.ras.ru/~duzhin/papers/>.
- [Ka] A. Kawauchi, *The invertibility problem on amphicheiral excellent knots*. Proc. Japan Acad. Ser. A Math. Sci. 55 (1979), no. 10, 399–402.
- [Kon] M. Kontsevich, *Vassiliev’s knot invariants*, Adv. in Soviet Math., **16** Part 2 (1993) 137–150.
- [L1] X.-S. Lin, *Finite type link invariants and the invertibility of links*, Math. Res. Letters 3 (1996), no. 3, pp. 405–417. Online at [arXiv:q-alg/9601019](http://arxiv.org/abs/q-alg/9601019).
- [L2] X.-S. Lin, *Finite type link-homotopy invariants*, l’Enseignement Mathématique 47 (2001), pp. 315–327. Online at [arXiv:math.GT/0012095](http://arxiv.org/abs/math.GT/0012095).
- [LM] T. Q. T. Le, J. Murakami, *The universal Vassiliev–Kontsevich invariant for framed oriented links*, Compositio Math. **102** (1996) 41–64.
- [Sta] T. Stanford, *Finite-type invariants of knots, links and graphs*. Topology, Vol. 35, no. 4, pp. 1027–1050.
- [Tro] H. F. Trotter, *Non-invertible knots exist*. Topology 2 (1964), 275–280.
- [VP] E. B. Vinberg and V. L. Popov, *Invariant theory*. In: Itogi Nauki Tekh., Ser. Sovrem. Probl. Mat., Fundam. Napravleniya 55 (1989), 137–314. English transl.: Encyclopaedia Math. Sci. vol 55, Springer, 1994.
- [Vog] P. Vogel, *Algebraic structures on modules of diagrams*, Institut de Mathématiques de Jussieu, Prépublication 32, August 1995, Revised in 1997, <http://www.math.jussieu.fr/~vogel/>.
- [W] H. Weyl, *The Classical Groups: Their Invariants and Representations*. Princeton University Press, 1997.

duzhin@pdmi.ras.ru